## **CYATHEACEAE**

**Alsophila costularis** Baker [= Cyathea chinensis Copel.]



Alsophila costularis Baker
Photo Courtesy: B.S. Kholia

Common English name: Tree fern.

Distribution: INDIA: Sikkim, Arunachal Pradesh, Mizoram (Lushai Hills) and Meghalaya (Khasia Hills); CHINA; NEPAL; BHUTAN; MYANMAR; THAILAND and VIETNAM.

Habitat: Grows at altitudes between 900 and 2300 m and also along forest fringes in open places and water channels.

Population status: Least Concern.

Description: Tree fern, 1.8-7.5 m tall with loose crown; trunk covered with thick black root fibres; rhizomes densely scaly, scales  $2.2-5 \times 1-1.5$  mm, dark brown. Stipes dark purple or stramineous-brown, smooth to warty or spinose to warty at base, warty to muricate above; pinnae rachis and costae clothed with crisp dark brown hairs on upper surface, pinnae  $50-75 \times 10-20$  cm, pinnules  $8-11 \times 1.5-2$  cm, deeply lobed; costules 3-4.5 mm apart; veins 7-10 pairs, usually forked. Sori yellow, round, raspberry like and naked, indusiate near costules; indusium attached only on side adjacent to costules, reflexed on maturity with abraded edge backing the costule; paraphyses shorter than sporangia; costules hairy throughout except near apex of pinnule on the lower surface.